9100200

<u>TO ALL TO WHOM THESE; PRESENTS; SHALL; COME;</u> The Board of Regents University of Nebraska and USBA-ARS Colherens, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED to be entitled to a certificate of plant variety protection under the LAW.

NOW, therefore, this certificate of plant variety protection is to grant UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, (THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT ETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. United States seed of this variety (1) shall be sold by variety name only as

OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.) Waived, except that this waiver shall not apply to breeder seed, n seed, labeling requirements, and blending limitations.)

'Rawhide'

In Testimony Wathereof, I have hereunto set my hand and caused the seal of the Wlant Variety Protection Office to be affixed at the City of Washington, D.C. 29th day of April the year of our Lord one thousand nine

hundred and ninety-four.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Expires 1/31/91

U.S. DEPARTMENT OF AGR AGRICULTURAL MARKETIN	Application is required in order to determine if a plant variety protection						
APPLICATION FOR PLANT VARIETY (Instructions on rec	certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).						
1. NAME OF APPLICANT(S) (as it is to appear on the Certificate) Board of Regents, University of Nebra.	o le o	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NO.	3. VARIETY NAME				
and USDA/ARS	Ska	NE 83498	Rawhide				
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)	····	PHONE (Include area code)	FOR OFFICIAL USE ONLY				
Lincoln, Nebraska 68583-0915		402-472-7211	PVPO NUMBER				
Washington, D.C. 20250		202-447-3656	9100200				
			13.1991				
6. GENUS AND SPECIES NAME 7.	FAMILY NAME (Botanie	cal)	Time				
Triticum aestivum L	Graminae	40	G A.M. P.M.				
8. CROP KIND NAME (Common Name)	9.	DATE OF DETERMINATION	F Filing and Examination Fee:				
Hard Red Winter Wheat	a)	July, 1983	s 2/50.				
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGANIZA	ATION (Corporation, part	nership, association, etc.)	R June 13,1991				
Corporation and U.S. Government Agency	7	b KHE	C Certificate Fee:				
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DA	TE OF INCORPORATION	\$ 250.00				
Nebraska and District of Columbia			E Avril 4 1994				
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SEL			5 17 1 p CCC 1, 1 / 7 / 1				
Dr. D. W. Nelson, Dean and Director		R. D. Plowman					
Agricultural Research Div., IANR-UNL Lincoln, Nebraska 68583-0704		A/ARS, Administration	Bldg., Room 302A				
Telephone: 402-472-2045	wası	nington, D.C. 20250	•				
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow I	MOTELLOTIONS	PHONE (Include area code):					
a. X Exhibit A, Origin and Breeding History of the Variety.	NSTRUCTIONS on rever	se)					
b. X Exhibit B, Novelty Statement.							
c. X Exhibit C, Objective Description of Variety.							
d. X Exhibit D, Additional Description of Variety.							
e X Exhibit E, Statement of the Basis of Applicant's Ownership.							
Seed Sample (2,500 viable untreated seeds). Date Seed Sar	mple mailed to Plant V	ariety Protection Office					
g. X Filing and Examination Fee (\$2,150) made payable to "Treasurer of the United States."							
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOLD E	BY VARIETY NAME ONLY	AS A CLASS OF CERTIFIED SEED? (See :	section 83(a) of the Plant Variety				
X YES (If "YES," answer items 16 and 17 below)		O," skip to item 18 below)	:				
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS?	i 17. IF "YES" TO	TEM 16, WHICH CLASSES OF PRODUCT	ION BEYOND BREEDER SEED?				
X YES NO	X YES NO X FOUNDATION X REGI						
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VARIET	TY IN THE U.S.?						
YES (If "YES," through Plant Variety Protection Act Patent Act. Give date:)							
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MARK		3.00					
X YES (If "YES," give names of countries and dates) United States, September, 1990							
∐ NO							
20. The applicant(s) declare(s) that a viable sample of basic seeds request in accordance with such regulations as may be applical	of this variety will ole.	be furnished with the application	and will be replenished upon				
The undersigned applicant(s) is (are) the owner(s) of this sexually reproduced novel plant variety, and believe(s) that the variety is distinct, uniform, and stable as required in section 41, and is entitled to protection under the provisions of section 42 of the Plant Variety Protection Act.							
Applicant(s) is (are) informed that false representation herein can jeopardize protection and result in penalties.							
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR T		DATE				
Daniel (2) Nolam)	Director	, Nebrusian station	4/12/91				
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR T		DATE				
	SEE ESTITION I	· 					
Administrator, ARS APR 2 9 1991							

FORM CSSD-470 (5-89) Edition of FORM LS-470, 3-86, is obsolete.

'Rawhide' (P. I. 518591) Hard Red Winter Wheat Application

Exhibit A. Origin and Breeding History:

Rawhide is a selection from the cross 'Warrior'*5/'Agent' //'Kavkaz' /4/'Parker'*4/Agent //'Beloterkovskaia 198' /'Lancer'/3/'Vona' made in 1977 by Dr. J. W. Schmidt. The F_1 generation was grown in the greenhouse in 1978. The F_2 and F_3 generations were grown in bulk at Mead, Nebraska in 1979 and 1980, respectively. Random heads were chosen from the F_3 and planted as head rows which were harvested in 1981. The F_3 -derived F_5 family was selected and harvested as a single observation plot in 1982. In 1983, Rawhide was grown at five locations in unreplicated trials in Nebraska. Rawhide was identified in these trials as NE83498. It has been tested in replicated trials from 1984 to the present. In addition, it has been tested in the Southern Regional Performance Nursery in 1989 and 1990. Rawhide was named and released on December 14, 1990 by the Nebraska Agricultural Experiment Station, South Dakota Experiment Station, and the Agricultural Research Service, U. S. Department of Agriculture. The initial allocation of Foundation seed to certified growers was made in September, 1990.

Rawhide will be maintained by the Nebraska Agriculture Experiment Station with the following classes: Breeder, Foundation, Registered, and Certified. Breeder seed of Rawhide will be maintained by roguing Breeder seed fields. The U.S. Department of Agriculture will not have seed for distribution.

Rawhide appears stable and uniform during seed increase. Less than 0.1% of the plants were rogued from Foundation and Breeder seed fields. It is expected that less than 0.1% (1:1000) variant plants (taller, 3 to 7 cm taller) may be encountered in subsequent generations.

Exhibit B. Novelty Statement

Rawhide is most similar to the hard red winter wheat cultivar Redland, but it can be distinguished by the following characteristics.

- 1. Rawhide is slightly shorter (25 mm) than Redland.
- 2. Rawhide is two days earlier than Redland.
- 3. Rawhide has about 1.8 lbs/bu heavier test weight.
- 4. In data provided by Dr. Don McVey of the USDA Cereal Rust Laboratories, Rawhide contains $\underline{Sr17}$ and $\underline{Sr24}$, is heterogeneous for $\underline{Sr31}$, whereas Redland contains genes $\underline{Sr5}$, $\underline{Sr17}$, and $\underline{Sr24}$.
- 5. Rawhide has hairs on the lower leaf auricles whereas Redland does not have hairy auricles.
- 6. Rawhide has a shorter (6.98 \pm 0.08 cm) and wider (11.1 \pm 0.2 mm) spike than Redland (7.96 \pm 0.14 cm and 9.7 \pm 0.1 mm)



Institute of Agriculture and Natural Resources

Department of Agronomy Crop, Range, Soil & Water, Weed Sciences 279 Plant Science • P.O. Box 830915 Lincoln, NE 68583-0915 PH. (402) 472-2811 • FAX (402) 472-7904 BITNET: AGRO001 @ UNLVM



October 25, 1993

Dr. A. A. Atchley Plant Variety Protection Office NAL Building, Room 500 10301 Baltimore Blvd. Beltsville, MD 20705-2351

Dear Alan:

I am responding to your letter of October 8, 1993 to Dr. D. W. Nelson concerning additional information needed on 'Rawhide' hard red winter wheat. Rawhide can be most clearly differentiated from '7833' (developed by HybriTech Seed International, Inc., but sold to AGSECO) by Rawhide being susceptible to wheat soilborne mosaic virus (WSBMV) and 7833 being resistant to WSBMV. In the Kansas State University 1992 summary of disease and insect reactions (enclosed), 7833 was scored as a 1 for resistance to WSBMV whereas Rawhide was scored as and 8. The scale is 1 for resistance and 9 for complete susceptibility. Resistance to WSBMV is important in eastern Kansas and Southeastern Nebraska, hence this is an important differentiating trait. Again using a 1 (resistant) to 9 (susceptible) scale, Rawhide is moderately resistant (4) to powdery mildew while 7833 is moderately susceptible (6). Similarly, Rawhide is moderately susceptible (6) to Hessian fly whereas 7833 is susceptible (8) to Hessian fly. I hope these traits will be sufficient to differentiate between Rawhide and 7833.

You also requested the number of generations over which Rawhide has been observed for stability and uniformity. Rawhide is stable and uniform over six generations of selfing and during the seed increase program. Finally you requested the characteristics which were bred for in this variety. I believe this is equivalent to asking for the criteria of selection. Our criteria for selection for each released hard red winter wheat cultivar are: a) adequate winterhardiness for propagation in Nebraska, b) resistance to <u>Puccinia graminis</u> (the causal organism of stem rust), c) agronomic performance equal or superior to commonly grown varieties, and d) acceptable end-use quality (in this case for bread making). These are the traits for which Rawhide was bred.

I hope this information will be satisfactory to conclude the PVP Application for Rawhide.

P. Stephen Baenziger

Eugene W. Price Professor and Head

CC

R. Hammons

D. Thompson

D. Nelson

Exh. A

7. Rawhide has a longer beak (7.63 \pm 0.33 mm) than Redland (1.49 \pm 0.07mm).

Exhibit C. See attached sheet.

Exhibit D.

Additional Description of the Variety:

Rawhide is an awned, white glumed cultivar. The foliage is blue-green with a waxy bloom at anthesis. The spike is middense and mostly strap. At maturity, the spike is inclined to nodding. The glume is midlong and midwide. The glume shoulder is narrow and oblique to square. The beak is acuminate and long with a range of medium to very long. The kernel is short, red colored, hard textured, and elliptical to ovate. The kernel has no collar, rounded cheeks, midsize germ, large brush, and a narrow and shallow crease.

Rawhide has been tested in Nebraska yield nurseries since 1984, in the Southern Regional Performance Nursery in 1988 and 1989 and in the Northern Regional Performance Nursery in 1989. In five years (23 location-years) of testing in the Nebraska Intrastate Nursery, Rawhide's yield (3420 kg/h) was similar to 'Siouxland' (3430 kg/h), less than 'Arapahoe' (3490 kg/h), and superior to 'Redland' (3300 kg/h). Yield, test weight, height, and anthesis

date will vary with the environment.

It is moderately resistant to the currently prevalent races of stem rust (incited by <u>Puccinia graminis</u> Pers. f. sp. <u>tritici</u> Eriks. and E. Henn.; contains <u>Sr17</u>, <u>Sr24</u>, and/or <u>Sr31</u>). It expresses the heterogeneous reaction to the Great Plains biotype of Hessian fly (<u>Mayetiola destructor</u> Say) which is believed to indicate the Marquillo-Kawvale genes for resistance. It is susceptible to leaf rust (incited by <u>P. recondita</u> Rob. ex. Desm. f. sp. <u>tritici</u> Eriks.), wheat soilborne mosaic virus, and wheat streak mosaic virus.

The overall bread making properties of Rawhide are average. Using composite samples from field trials in Nebraska, Rawhide is similar to Arapahoe and Scout 66 in wheat and flour protein content and higher than Centurk 78. Rawhide is also similar to Arapahoe, less than Scout 66, and greater than Centurk 78 for milling extraction. Baking absorption was similar for Rawhide, Arapahoe, Centurk 78, and Scout 66. Rawhide has strong mixing characteristics as determined by the mixograph. The loaf volume of Rawhide was less than Arapahoe and Scout 66, and was similar to Centurk 78. Internal and external loaf characteristics were acceptable (see attachement). The kernels of Rawhide have been classified, using the existing standards, by the Federal Grain Inspection Service as being hard red winter wheat.

Exhibit E.

Statement of the Basis of the Applicant's Ownership

The University of Nebraska and the USDA/ARS are the applicants for protection in the case of Rawhide hard red winter wheat being:

a) Rawhide is a product of the cooperative state-federal breeding program location in the Agricultural Research Division (ARD), University of Nebraska. Drs. P. Stephen Baenziger, John W. Schmidt, and Paul J. Mattern;

- and Drs. C. James Peterson and Virgil A. Johnson, regular employees of the Nebraska ARD (Department of Agronomy) and the USDA/ARS (stationed and functioning also as staff members in the Department of Agronomy), respectively, have bred the named cultivar for and within these incorporated institutions.
- b) By established policy, release of cultivars developed by the Nebraska ARD is the responsibility of the Nebraska ARD as the agency providing staff, funds, and facilities for the breeding program.

U, S, DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
LIVESTOCK, MEAT, GRAIN AND SEED DIVISION
BELTSVILLE, MARYLAND 20786
OBJECTIVE DESCRIPTION OF VARIETY

EXHIBIT C (Whoat)

INSTRUCTIONS: See Reverse. WHEAT (TRITICUM SP	'P.)
Board of Regents, University of Nebraska and	FOR OFFICIAL USE ONLY
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	9100200
Lincoln, NE 68583-0745	VARIETY NAME OR TEMPORARY DESIGNATION
Washington, DC 20750	
Place the appropriate number that describes the varietal character of this varietal energy of this varietal character of this var	iety in the boxes below.
1. KIND:	or ress or y or ress.
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH	6 = POULARD 7 = CLUB
2. TYPE,	OFT 3 = OTHER (Specify)
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify) 2 2 = H	
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO: NA	The same of the sa
FIRST FLOWERING	LAST FLOWERING
4. MATURITY (50% Flowering):	
0 2 NO. OF DAYS EARLIER THAN	ARTHUR 2 = SCOUT 3 = CHRIS
NO. OF DAYS LATER THAN	EMHI 5 = NUGAINES 6 = LEEDS Redland
PLANT HEIGHT (From soil level to top of head):	REGIEIM
0 8 4 cm. High	
CM. TALLER THAN	Redland
2. 5 CM. SHORTER THAN	RTHUR 2 = SCOUT 3 = CHRIS
PLANT COLOR AT BOOTING (See reverse): 7. ANTHER	EMHI 5 = NUGAINES 6 = LEEDS
3 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	LLOW 2 = PURPLE
STEM:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT 2 Waxy bl	oom: 1 = ABSENT 2 = PRESENT
Hairiness of last internode of rachis: = ABSENT 2 = PRESENT Internod	les: 1 = HOLLOW 2 = SOLID
	1. INTERNODE LENGTH BETWEEN FLAG LEAF ND LEAF BELOW
AURICLES:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT 2 Hairings	s: 1 = ABSENT 2 = PRESENT
LEAF:	
Flag leaf at 1 = ERECT 2 = RECURVED 2 Flag leaf stage: 3 = OTHER (Specify): 2 Flag leaf	if: 1 = NOT TWISTED 2 = TWISTED
Hairs of first last about 1 - annual	oom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
0 MM. LEAF WIOTH (First loss below flag loss) 1 7 CM	· LEAF LENGTH (First leaf below flag loaf):
RM LMGS 470-5 (6-82) (Formerly Form LPGS 470.6 /2 70)	

11. HEAD:				
3 Density: 1 = LAX 2 = DENSE 3= Mid	dense 2		NG 2 = STRAP 3 = (Specify)	CLAVATE
4 Awnedness: 1 = AWNLESS 2 = APICALLY	AWNLETED 3 = AWNLE	red 4 = AWNE	· , , , , , , , , , , , , , , , , , , ,	4.4
Color at maturity: 5 = BROWN 6 = BLAC	3 = PINK 4 = RED K 7 = OTHER (Specify)	·	No. of the second	-
7. 0 CM. LENGTH	1	1 MM. WIDTH		
12. GLUMES AT MATURITY: 2 Length: 1 = SHORT (CA. 7 mm.) 2 = MEDI 3 = LONG (CA. 9 mm.)	UM (CA. 8 mm.)	/idth: 1 = NARROW 3 = WIDE (C.	(CA. 3 mm.) 2 = MEDII A. 4 mm.)	JM (CA. 3.5 mm.)
Shoulder 1 = WANTING 2 = OBLIQUE 3 shape: 4 = SQUARE 5 = ELEVATED		eak: l=08TUSE	2 = ACUTE 3 = AC	UMINATE
13. COLEOPTILE COLOR:	14. SEE	DLING ANTHOCY	ANIN:	
1 i = white 2 = RED 3 = PURPLE		= ABSENT 2	= PRESENT	
15. JUVENILE PLANT GROWTH HABIT:				
1 1 = PROSTRATE 2 = SEMI-ERECT	3 = ERECT	· · · · · · · · · · · · · · · · · · ·		
16. SEED:				
With a tendency to ovate 3 = ELL	PTICAL 1 C	heek: 1 = ROUNDI	ED 2 = ANGULAR	
2 Brush: 1 = SHORT 2 = MEDIUM 3 = LC	NG 1 E	tush: = NOT CC	DLLARED 2 = COLLAR	ED
Phenol reaction 1 = IVORY 2 = FAWN 5 = BLA				
Color: 1 = WHITE 2 = AMBER 3 = RED	4 = PURPLE 5 = OTI	HER (Specify)		· · · · · · · · · · · · · · · · · · ·
5. 8 MM. LENGTH 2. 9 MM. WI	2 [GM. PER 1000	SEEDS	
17. SEED CREASE:				
Width: 1 = 60% OR LESS OF KERNEL WINO	(A' 1 D	epth: 1 = 20% OF	LESS OF KERNEL SCOL	ידי
2 = 80% OR LESS OF KERNEL 'CHRIS		2 = 35% OR	LESS OF KERNEL 'CHRI	5'
3 = NEARLY AS WIDE AS KERNEL 'LE	мни	3 = 50% OR	LESS OF KERNEL 'LEMP	1(*
18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 =	Resistant)			
2 STEM RUST TMNK 1 LEAF RUS (Races)		TRIPE RUST Races)	0 LOOSE SMU	τ
0 POWDERY MILDEW 0 BUNT	· 125 o	THER (Specify)		·
19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 =	Resistant) 3= Moderate	ly Resistan	t	•
O SAWFLY O APHID (Byd		REEN BUG		AF BEETLE
OTHER (Specify)	SESSIAN FLY	0 A	0 в	<u>0</u> c
	RACES: 0 D	Ο ε	0 F	0 G
20. INDICATE WHICH VARIETY MOST CLOSELY RE	SEMBLES THAT SUBMITTED):		
CHARACTER NAME OF V		ARACTER	NAME OF VAR	ETY
Plant tillering Red1and		eed size	Centurk 78 🔍	
Leaf size Redland	S	eed shape	Arapahoe 🖄	
Leaf color Cody	Coleo	otile elongation	Arapahoe/	100 July X
Leof corriage Redland	Seedli	ng pigmentation	Siouxland /	Les Alle
GENERAL: The following publications may be used:	and the second of the second o	i .	,	. 977

(b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

F COLOR: Nickerson's or say the Association of Official Seed Analysts.

- LEAF COLOR: Nickerson's or any recognized color fan should be used to determine the leaf color of the described variety.

er e	Test Weight (lb/bu)					
	1989	1990	Average			
	(11)*	(12)	(23)			
Rawhide	57.5	58.3	57.9			
Redland	55.8	56.3	56.1			

* (Number of locations)

Table 2.

	Protein (%)					
	1989	1990	Average			
	(11)	(12)	(23)			
Rawhide	13.7	12.2	12.9			
Redland	12.9	11.4	12.1			

* (Number of locations)

Table 3. Nebraska Interstate Nursery.

	Yield (bu/a)							
	1986	1987	1988	1989	1990	Average		
	(5)*	(5)	(4)	(4)	(5)	(23)		
Arapahoe	54.3	58.3	51.1	42.0	53.9	51.9		
Colt	44.8	55.7	44.0	37.4	49.1	46.2		
Cody	55.4	53.8	49.7	39.3	50.2	49.7		
Redland	47.7	56.8	48.7	40.4	51.5	49.0		
Siouxland	52.2	55.0	54.5	42.4	51.0	51.0		
Rawhide	46.8	57.9	55.6	42.4	51.6	50.9		

* (Number of locations)

Table 4. Comparison of Rawhide and Redland for spike and coleoptile characteristics.

	Rawhide	Red1 and
Beak length (mm)	7.63±0.32	1.49±0.07
Glume length (mm)	7.90 ± 0.07	7.10+0.06
Glume width (mm)	3.44 ± 0.05	3.30 ± 0.04
Spike length (cm)	6.98 + 0.08	7.96+0.14
Spike width (mm)	11.12 ± 0.20	9.72 + 0.14
Spike density (cm)	3.67 ± 0.05	3.81 ± 0.06
Coleoptile length (mm)	40.54 ± 1.07	37.18 ± 0.34

RAWHIDE

HARD RED WINTER WHEAT

MILLING AND BAKING PROPERTIES

VARIETY	WHT	FLR	FLR	MILL	MIXOG	RAPH			BA	KING		
(YEARS TESTED)	PRO %	PRO %	ASH %	YLD %	TIME min	TOL	ABS %	MIX min	CC	EXT	GRN	TEX
RAWHIDE (1983-89)	12.3	11.3	0.41	72.4	5.6	4+	61.4	5.9	861	G-	G+	G+
ARAPAHOE (1983-89)	12.5	11.3	0.45	72.1	4.9	4	61.3	5.2	883	G	G+	G+
SCOUT 66 (1983-89)	12.2	11.1	0.39	73.5	3.2	3+	61.9	3.2	876	G	G	G
CENTURK 78 (1983-85, 87,88)	11.8	10.6	0.41	71.0	5.5	5-	61.2	5.9	858	G-	VG	VG-

The data represent composite averages for the years as indicated and were reported by the Wheat Quality Laboratory, University of Nebraska, Lincoln. Rawhide was similar to Arapahoe and Scout 66 in wheat and flour protein content and higher than Centurk 78. The milling extraction for Rawhide was similar to Arapahoe, less than Scout 66, and greater than Centurk 78. The flour ash level was The dough rheological properties of Rawhide, acceptable. determined with the mixograph, indicated strong mixing properties that are presently desired by the milling and baking industry. The baking absorption of the experimental variety was similar to the named varieties. The mixing time is longer than Arapahoe and Scout 66 and similar to Centurk 78. The loaf volume was less than Arapahoe and Scout 66 and similar to Centurk 78. External and internal characteristics of the baked bread were acceptable.

CONCLUSIONS

Wheat Protein Content	Average
Milling Properties	Average
Mixing Properties	Strong
Baking Characteristics	Acceptable

Prepared: January 1991 David R. Shelton

Plant Variety Protection Office NAL Building, Rm. 500 10301 Baltimore Blvd. Beltsville, MD 20705-2351

United States Department of Agriculture

PLANT VARIETY PROTECTION OFFICE

Gentlemen:

Subject: Application No.

9100200

Variety and Kind:

"Rawhide" Hard red winter wheat

As provided in section 83(a) of the Plant Variety Protection Act, 7 U.S.C. 2321, we request that the Certificate on the above variety be issued with a notation on the Certificate that the right to exclude others from selling, offering for sale, reproducing, importing or exporting the variety covered by this Certificate, or using it in producing a hybrid or different variety is <u>waived</u>, except that this waiver shall not apply to breeders seed, foundation seed, labeling requirements, and blending limitations.

It has been agreed that the Certificate should be issued in the name(s) of:

Board of Regents, University of Nebraska and USDA/ARS

June 28, 1991

(Date)

Haule W Melson

(Signature)
Director, Nebraska Agricultural Experiment Station